

International Journal of Modern Physics D
 © World Scientific Publishing Company

DOES CASSINI ALLOW TO MEASURE RELATIVISTIC ORBITAL EFFECTS IN THE SATURNIAN SYSTEM OF SATELLITES?

LORENZO IORIO

*Viale Unità di Italia 68, 70125, Bari (BA), Italy
 lorenzo.iorio@libero.it*

Received Day Month Year

Revised Day Month Year

Communicated by Managing Editor

In this paper we address the following question: do the recent advances in the orbit determination of the major natural satellites of Saturn obtained with the analysis of the first data sets from the Cassini mission allow to detect the general relativistic gravitoelectric orbital precessions of the mean longitudes of such moons? The answer is still negative. The present-day down-track accuracy is adequate for Mimas, Enceladus, Thetys, Dione, Rhea and Titan and inadequate for Hyperion, Iapetus and Phoebe. Instead, the size of the systematic errors induced by the mismodelling in the key parameters of the Saturnian gravitational field like the even zonal harmonics J_ℓ are larger than the relativistic down-track shifts by about one order of magnitude, mainly for the inner satellites like Mimas, Enceladus, Thetys, Dione, Rhea, Titan and Hyperion. Iapetus and Phoebe are not sensibly affected by such kind of perturbations. Moreover, the bias due to the uncertainty in Saturn's GM is larger than the relativistic down-track effects for all such moons up to two orders of magnitude (Phoebe). Thus, it would be impossible to separately analyze the mean longitudes of each satellite. Proposed linear combinations of the satellites' mean longitudes would allow to cancel out the impact of the mismodelling in the low-degree even zonal harmonics and GM , but the combined down-track errors would be larger than the combined relativistic signatures by a factor 10^3 .

Keywords: Cassini spacecraft; general relativistic orbit precessions; Saturnian system of natural satellites

1. Introduction

A satisfactorily empirical corroboration of a fundamental theory requires that as many independent experiments as possible are conducted by different scientists in different laboratories. Now, general relativity is difficult to test, especially in the weak-field and slow-motion approximation, valid, e.g., in our Solar System, both because the relativistic effects are very small and the competing classical signals are often quite larger. Until now, Solar System tests of general relativity accurate to better than 1% have been performed by many independent groups only in the gravitational field of the Sun by checking the effects induced by the gravitoelectric

Schwarzschild^a part of the space-time metric on the propagation of electromagnetic waves and planetary orbital motions⁴. It is as if many independent experiments aimed to measure fundamental physical effects were conducted always in the same laboratory. Thus, it is worthwhile to try to use different laboratories, i.e. other gravitational fields, to perform such tests, even if their outcomes should be less accurate than those conducted in the Sun's field. In principle, the best candidates other than the Sun are the giant planets of the Solar System like Jupiter and Saturn. To date, the only investigations of this kind are due to Hiscock and Lindblom⁵, who preliminarily analyzed the possibility of measuring the Einstein pericentre precessions⁶ of some of the natural satellites of Jupiter and Saturn, and to Iorio and Lainey⁷ who investigated the measurability of the Lense-Thirring precessions in the system of the galilean satellites of Jupiter with modern data sets^b.

The aim of this paper is to investigate if the recent improvements in the ephemerides of the major Saturnian satellites (see Table 1 for their orbital parameters) obtained from the first Cassini data allows to detect at least the general relativistic gravitoelectric precessions of their orbits.

Table 1. Orbital parameters of the major Saturnian satellites: a is the semimajor axis, e is the eccentricity and i is the inclination, referred to the local Laplace planes (http://ssd.jpl.nasa.gov/?sat_elem).

Satellite	a (km)	e	i (deg)
Mimas	185540	0.0196	1.572
Enceladus	238040	0.0047	0.009
Thetys	294670	0.0001	1.091
Dione	377420	0.0022	0.028
Rhea	527070	0.0010	0.331
Titan	1221870	0.0288	0.280
Hyperion	1500880	0.0274	0.630
Iapetus	3560840	0.0283	7.489
Phoebe	12947780	0.1635	175.986

2. The relativistic effects investigated

The Einstein pericentre precession is undoubtedly the most famous relativistic orbital effect, but it is not the only one. Indeed, also the mean anomaly $\mathcal{M}$ undergoes

^aIn regard to the much smaller gravitomagnetic Lense-Thirring precessions¹ of the planetary orbits, they lie just at the edge of the present-day accuracy in planetary ephemerides and have recently been found in agreement with the latest measurements, although the errors are still large². A 6% test performed in the gravitational field of Mars with the MGS spacecraft has recently been reported in Ref. 3.

^bIn their original papers Lense and Thirring¹ proposed to use some of the satellites of Jupiter and Saturn.

Does Cassini allow to measure relativistic orbital effects in the Saturnian system of satellites? 3

a secular advance which is even larger than that experienced by the pericentre. For small eccentricities it is⁸

$$\dot{\mathcal{M}}_{\text{GE}} \sim -\frac{9nGM}{c^2 a \sqrt{1-e^2}}, \quad (1)$$

where G is the Newtonian constant of gravitation, M is the mass of the central body which acts as source of the gravitational field, a , e and $n = \sqrt{GM/a^3}$ are the semimajor axis, the eccentricity and the Keplerian mean motion, respectively, of the satellite's orbit. Eq. (1), which yields a rate of -130 arcseconds per century for Mercury, has been obtained by using the standard isotropic radial coordinate r related to the Schwarzschild coordinate r' by $r' = r(1 + GM/2c^2r)^2$. The validity of Eq. (1) has also been numerically checked by integrating over 200 years the Jet Propulsion Laboratory (JPL) equations of motion of all the planets of the Solar System, which are written in terms of just the standard isotropic radial coordinate, with and without the gravitoelectric $1/c^2$ terms in the dynamical force models⁹ in order to single out just the post-Newtonian gravitoelectric effects (E.M. Standish, private communication, 2004). The obtained precessions fully agree with those obtainable from Eq. (1).

As a consequence, the mean longitude^c $\lambda = \omega + \Omega + \mathcal{M}$, which is one of the non-singular orbital elements used for orbits with small eccentricities and inclinations like those of most of the Solar System major bodies as just the Saturnian satellites (see Table 1), experiences a secular advance

$$\dot{\lambda}_{\text{GE}} \sim -\frac{6nGM}{c^2 a}. \quad (2)$$

It is twice the pericentre advance. On the other hand, because of $\mathcal{M}$ in the definition of λ the systematic uncertainty in the Keplerian mean motion n must also be accounted for if a secular rate must be extracted from the analysis of such an orbital element.

3. The possibilities offered by the Saturnian system

Is it possible to measure the gravitoelectric orbital precessions of the natural satellites of Saturn, in view of the recent refinements of their ephemerides obtained by analyzing the first data from the Cassini spacecraft? In this Section we will address this problem in detail. To this aim, we must confront the magnitude of the relativistic effects of interest with the major systematic errors induced by classical forces having the same signatures (Section 3.1), and with the currently available measurement accuracies (Section 3.2).

3.1. The reduction of the impact of the even zonal harmonics

In regard to the first issue, a major source of systematic bias is represented by the even zonal harmonic coefficients J_ℓ , $\ell = 2, 4, 6, \dots$ of the multipolar expansion of the

^cHere ω and Ω are the argument of pericentre and the longitude of the ascending node, respectively.

Saturn's gravitational potential. Indeed, they induce secular precessions on λ

$$\dot{\lambda}_{\text{even zonals}} = \sum_{\ell \geq 2} \dot{\lambda}_{\ell} J_{\ell}, \quad (3)$$

where the coefficients $\dot{\lambda}_{\ell}$ are expressed in terms of the satellite's orbital elements a, e, i and the mass and the radius of the central body. Such classical advances must be accurately modelled because they are much larger than the relativistic ones. Even the latest determinations of the Saturnian gravity field from some Cassini data sets¹⁰ show that the currently available models of the even zonal harmonics do not reach the required accuracy to allow for a measurement of the Einstein precessions. This fact is clearly shown by Table 2. Indeed, we can note that for

Table 2. Mean longitudes of the major Saturnian satellites: general relativistic gravitoelectric secular precessions $\dot{\lambda}_{\text{GE}}$, in $'' \text{ cy}^{-1}$, mismodelled classical secular precessions $\delta\dot{\lambda}_{J_2}, \delta\dot{\lambda}_{J_4}$, in $'' \text{ cy}^{-1}$, due to the uncertainties in the even zonal harmonics J_2, J_4 , systematic errors δn_{GM} , in $'' \text{ cy}^{-1}$, in the Keplerian mean motions n due to the uncertainty in Saturn's GM . The values $\delta(GM) = 1.2 \text{ km}^3 \text{ s}^{-2}$, $\delta J_2 = 0.4 \times 10^{-6}$, $\delta J_4 = 3 \times 10^{-6}$ have been used.

Satellite	$\dot{\lambda}_{\text{GE}}$	$\delta\dot{\lambda}_{J_2}$	$\delta\dot{\lambda}_{J_4}$	δn_{GM}
Mimas	-684.670	5926.552	5465.353	793.461
Enceladus	-367.207	2478.985	1390.913	546.018
Thetys	-215.374	1173.870	429.231	396.440
Dione	-116.003	493.910	110.229	273.492
Rhea	-50.334	153.451	17.558	165.721
Titan	-6.152	8.101	0.172	46.950
Hyperion	-3.679	3.942	0.055	34.487
Iapetus	-0.424	0.186	$\mathcal{O}(10^{-4})$	9.437
Phoebe	-0.016	0.002	$\mathcal{O}(10^{-7})$	1.361

Mimas, Enceladus and Thetys the mismodelled precessions due to J_2 and J_4 are larger than the relativistic rates by about one order of magnitude. For Dione, Rhea, Titan and Hyperion the bias due to J_4 is smaller than the relativistic signal. For Iapetus and Phoebe the even zonal harmonics of Saturn do not represent a problem. In addition to the bias due to the $\{J_{\ell}\}$, for the mean longitude there is also the systematic error $\delta n_{GM} = \delta(GM)/\sqrt{4GMa^3}$ induced on the Keplerian mean motion n by the uncertainty in the Saturn's GM which, as can be seen, is larger than the relativistic shifts for all the satellites.

A way to overcome the problem of the impact of the even zonals is represented by the linear combination approach^{8,11} summarized in the following. Let us write down the expressions of the measured residuals^d of the mean longitudes $\delta\dot{\lambda}_{\text{meas}}$ of

^dThe Keplerian orbital elements are not directly observable, so here we are using the words 'measured residuals' in an improper sense. We mean, instead, a set of solve-for parameters of a suitable

Does Cassini allow to measure relativistic orbital effects in the Saturnian system of satellites? 5

N chosen satellites in terms of the mismodelled Keplerian mean motions, of the classical precessions due to the mismodelled parts of the even zonal harmonics, and of the relativistic precessions, assumed as totally unmodelled features of motion

$$\delta \dot{\lambda}_{\text{meas}}^{(j)} = n_{.GM} \delta(GM) + \sum_{\ell \geq 2}^{2(N-2)} \dot{\lambda}_{\ell}^{(j)} \delta J_{\ell} + \dot{\lambda}_{\text{GE}}^{(j)} \xi, \quad j = 1, 2..N, \quad (4)$$

where $\dot{\lambda}_{\ell}$ are the coefficients of the classical precessions of degree ℓ of the mean longitude and

$$n_{.GM} = \frac{1}{\sqrt{4GMa^3}}. \quad (5)$$

If we look at Eq. (4) as a inhomogeneous algebraic system of N linear equations in the N unknowns $\{\delta(GM); \delta J_{\ell}; \xi\}$ and solve it for ξ , we get a linear combination of the satellites' mean longitudes residuals which, by construction, is independent of $\delta(GM)$ and the first $N - 2$ even zonal harmonics, and is sensitive just to the relativistic signatures and to the perturbations of the remaining, uncanceled even zonal harmonics. The cancellation of the low-degree even zonals is important also because in this way one avoids any possible a priori 'imprint' of the relativistic effects themselves via such spherical harmonics. Indeed, the Saturnian gravity field models are obtained as least-square solutions using all the available data from the spacecraft encounters and the satellites' motions around Saturn, so that the relativistic effects themselves are in some way included in the solved-for parameters like $\{J_{\ell}\}$.

A possible combination which cancels out δJ_2 , δJ_4 and $\delta(GM)$ is

$$\xi = \frac{\delta \dot{\lambda}_{\text{meas}}^{\text{Ph}} + k_1 \delta \dot{\lambda}_{\text{meas}}^{\text{Ia}} + k_2 \delta \dot{\lambda}_{\text{meas}}^{\text{Hy}} + k_3 \delta \dot{\lambda}_{\text{meas}}^{\text{Ti}}}{\dot{\lambda}_{\text{GE}}^{\text{Ph}} + k_1 \dot{\lambda}_{\text{GE}}^{\text{Ia}} + k_2 \dot{\lambda}_{\text{GE}}^{\text{Hy}} + k_3 \dot{\lambda}_{\text{GE}}^{\text{Ti}}}, \quad (6)$$

with

$$\left\{ \begin{array}{l} k_1 = -0.1296, \\ k_2 = 0.0224, \\ k_3 = -0.0067, \\ \dot{\lambda}_{\text{GE}}^{\text{Ph}} + k_1 \dot{\lambda}_{\text{GE}}^{\text{Ia}} + k_2 \dot{\lambda}_{\text{GE}}^{\text{Hy}} + k_3 \dot{\lambda}_{\text{GE}}^{\text{Ti}} = 0.0237 \text{ ''cy}^{-1} \end{array} \right. \quad (7)$$

As can be argued from Table 2, the impact of the remaining uncanceled even zonal harmonics J_6, J_8 is negligible.

least-square solution to be obtained by contrasting all the available observational data to a set of dynamical force models in which all the $1/c^2$ terms have purposely been switched off.

3.2. The present-day orbital accuracy

In regard to the feasibility of such a measurement, it is useful to consider the down-track shifts $\Delta\tau \sim a\Delta\lambda$ and the present-day accuracy according to the latest Saturnian ephemerides SAT240¹² over 81 years. The results are in Table 3. It can

Table 3. Major Saturnian satellites: gravitoelectric down-track shifts, in km, and present-day orbital accuracy, in km, over a time span of 81 years. See also on the Internet http://ssd.jpl.nasa.gov/?sat_ephem.

Satellite	$\Delta\tau_{\text{GE}}$ (km)	$\delta(\Delta\tau)$ (km)
Mimas	-498	50
Enceladus	-343	20
Thetys	-249	50
Dione	-171	20
Rhea	-104	20
Titan	-29	10
Hyperion	-21	100
Iapetus	-6	20
Phoebe	-0.8	50

be noted that the situation is presently not favorable just for the satellites used for the combination of Eq. (6). This fact reflects on Eq. (6) itself: indeed the total uncertainty calculated by summing in quadrature the errors $\delta(\Delta\tau)$ of Table 3 with the coefficients (7)

$$\left. \frac{\delta\xi}{\xi} \right|_{\text{meas}} = \frac{\sqrt{[\delta(\Delta\tau)^{\text{Ph}}]^2 + [k_1\delta(\Delta\tau)^{\text{Ia}}]^2 + [k_2\delta(\Delta\tau)^{\text{Hy}}]^2 + [k_3\delta(\Delta\tau)^{\text{Ti}}]^2}}{\Delta\tau_{\text{GE}}^{\text{Ph}} + k_1\Delta\tau_{\text{GE}}^{\text{Ia}} + k_2\Delta\tau_{\text{GE}}^{\text{Hy}} + k_3\Delta\tau_{\text{GE}}^{\text{Ti}}} \quad (8)$$

is $\sim 10^3$ times larger than the gravitoelectric shift over 81 years.

About the use of the inner satellites, for which the relativistic shifts $\Delta\tau_{\text{GE}}$ are larger than the errors $\delta(\Delta\tau)$, Table 2 shows that it is not possible to use their mean longitudes without combining them because both the even zonal mismodelled shifts and the bias of n are quite larger. On the other hand, it turns out that also the linear combination approach does not yield good results. Indeed, the root-sum-square of the errors of a combination with Enceladus, Dione, Rhea and Titan is 10^3 times larger of the relativistic combined shifts.

4. Discussion and conclusions

In regard to the measurability of the general relativistic gravitoelectric orbit advances in the Saturn's system of natural satellites, it turns out that the present-day improvements in their ephemerides by the Cassini spacecraft are not yet sufficient to detect such post-Newtonian effects.

As a general rule, in regard to the single satellites, an about one order of magnitude improvement in the knowledge of the parameters of the Saturnian gravity field

Does Cassini allow to measure relativistic orbital effects in the Saturnian system of satellites? 7

like the even zonal harmonics J_ℓ and GM would be required to make the competing classical bias smaller than the relativistic precessions. This would make possible to successfully analyze separately the mean longitudes of each of Mimas, Enceladus, Thetys, Dione, Rhea and Titan because the present-day down-track errors are already smaller than the relativistic shifts. The use of the mean longitudes of Hyperion, Iapetus and Phoebe, for which the Saturnian even zonals represent a relatively less important problem, is strictly related to a one-two orders of magnitude improvements in the down-track parts of their orbits.

The linear combination approach yields, in principle, good results in reducing the impact of the mismodelling in the even zonals and GM . The main problem with such strategy relies in the down-track orbital accuracy. Indeed, the down-track errors weighted by their combination's coefficients must be summed in quadrature, while the weighted relativistic shifts are combined with their proper own signs, so that with the present-day precision the overall error amounts to 10^3 .

References

1. J. Lense and H. Thirring H., *Phys. Z.* **19**, 156 (1918). Translated and discussed by B. Mashhoon, F.W. Hehl and D.S. Theiss, *Gen. Relativ. Grav.* **16**, 711 (1984). Reprinted in R.J. Ruffini and C. Sigismondi (eds), *Nonlinear Gravitodynamics*. (World Scientific, Singapore, 2003). p. 349.
2. L. Iorio gr-qc/0507041 (2005a).
3. L. Iorio, *Class. Quantum Grav.* , **23**, 5451 (2006).
4. C.M. Will, *Living Rev. Relativity* **9**, 3 (2006). (www.livingreviews.org/lrr-2006-3)
5. W.A. Hiscock and L. Lindblom, *Astron. J.* **231**, 224 (1979).
6. A. Einstein, *Sitzber. Kön. Preuss. Akad. Berlin.* **47**, 831 (1915).
7. L. Iorio and V. Lainey, *Int. J. Mod. Phys. D* **14**, 2039 (2005).
8. L. Iorio, *Astron. Astroph.* **433**, 385 (2005b)
9. F. Estabrook, *JPL IOM, Section 328*, 1 (1971).
10. R.A. Jacobson *et al.*, *American Astronomical Society, DPS meeting #37, #47.11* (2005).
11. L. Iorio, *New Astron.* **10**, 603 (2005c)
12. R.A. Jacobson, *SAT240, JPL Saturn Ephemerides* (2006).